IL LATTOSIO È LO ZUCCHERO NATURALMENTE CONTENUTO NEL LATTE E NEI PRODOTTI DERIVATI

Latte

6 g di lattosio
(1 bicchiere = 125 mL)

Yogurt

5 g di lattosio
(1 vasetto = 125 g)

Formaggi stagionati

Tracce di lattosio (50 g)
(1tagliere)

L'ENZIMA LATTASI SEPARA IL LATTOSIO IN DUE ZUCCHERI: GLUCOSIO E GALATTOSIO

LA RIDUZIONE FISIOLOGICA DELL'ATTIVITÀ DELLA LATTASI COMPORTA UNA MINORE DIGESTIONE DEL LATTOSIO

QUANDO LA DIFFICOLTÀ A DIGERIRE IL LATTOSIO COMPORTA SINTOMI SPECIFICI SI PARLA DI INTOLLERANZA AL LATTOSIO

L'INTOLLERANZA AL LATTOSIO È MOLTO MENO DIFFUSA

COME SI PUÒ ACCERTARE L'INTOLLERANZA AL LATTOSIO?

E' NECESSARIO RIVOLGERSI AL PROPRIO MEDICO: LA DIAGNOSI DI INTOLLERANZA AL LATTOSIO PUÒ ESSERE POSTA SOLTANTO DAL MEDICO CHE, SE OPPORTUNO, POTRÀ DECIDERE DI PRESCRIVERE IL TEST DEL RESPIRO (HYDROGEN BREATH TEST)

ATTENZIONE ALLE DIAGNOSI FAI-DA-TE

1

DOPO
50%
AGGIUNGERE:
DEI CASI

La diagnosi fai da te spesso si associa a sintomi diversi e ad una scarsa qualità della vita complessiva

2

Questi sintomi non sono specifici dell'intolleranza al lattosio

Fattori psicologici
(ansia, stress, depressione)

Patologie intestinali
(morbo di Crohn e colite ulcerosa)

→ Flatulenza
→ Diarrea
→ Gonfiore

3

Intolleranza al lattosio ≠ Allergia alle proteine del latte vaccino

NON È NECESSARIO EVITARE IL LATTE E I DERIVATI



Riduzione o eliminazione di latte e derivati

Possibile rischio di carenza nutrizionale

E' SUFFICIENTE RIVEDERE LE PROPRIE ABITUDINI ALIMENTARI

Evitare gli eccessi è la regola principale, valida anche per chi non digerisce il lattosio o è intollerante

I fermenti vivi dello yogurt favoriscono la digestione del lattosio

Alcuni formaggi stagionati sono a basso/nullo contenuto di lattosio

Gli alimenti che contengono lattosio possono essere consumati in quantità limitate nel corso della giornata

EFSA LO HA CONFERMATO: I FERMENTI VIVI DELLO YOGURT MIGLIORANO LA DIGESTIONE DEL LATTOSIO

ALTRE ISTITUZIONI E SOCIETÀ MEDICO-SCIENTIFICHE CONFERMANO CHE LO YOGURT PUÒ ESSERE CONSUMATO ANCHE DA CHI HA DIFFICOLTÀ A DIGERIRE IL LATTOSIO

VITAMINE E MINERALI SONO ESSENZIALI...

Lo yogurt è un **alimento che può essere consumato da tutti** nell'ambito di **un'alimentazione varia ed equilibrata**.

Bibliografia:
Adolfsson O and al., Yogurt and gut function. Am J Clin Nutr 2004;80:245-56.
Bailey RK and al., Lactose intolerance and health disparities among African Americans and Hispanic Americans: an updated consensus statement. J Natl Med Assoc 2013;105:112-27.
Casellas F and al., Lactose intolerance: an overview. Rev Esp Enferm Dig (Madrid) 2008;100:551-8, 559-64.
Casellas F, Aparici A, Perez M, J, Rodriguez P, (2015) Prevalence of lactose intolerance impairs health-related quality of life. European journal of clinical nutrition
Commission CA. Codex standard for fermented milks. Food and Agriculture Organization United Nation Rome 2003;5:1.
Commission CA. Codex standard for fermented milks. Food and Agriculture Organization United Nation Rome 2003;5:1.
Corrigan CJ and al., Fate of milk lactose in allergic children with cow's milk allergy. Pediatr Allergy Immunol 2002;13:70-4.
Elisa Pesce on Dietary Products N. Allergen, Scientific Opinion on the substantiation of health claims related to live yoghurt cultures and improved lactose digestion (ID 1143, 2976) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. EFSA Journal 2010;8:1763-1763.
Fassio F and al., Lactose maldigestion, malabsorption, and intolerance: a comprehensive review with a focus on current management and future perspectives. Nutrients 2018;10:1599.
Guarner F, Milk and dairy products: a unique micronutrient combination. J Am Coll Nutr 2011;30:150-9.
Guarner F, Milk and dairy products: a unique micronutrient combination. J Am Coll Nutr 2011;30:150-9.
Lukito W and al., From "lactose intolerance" to "lactose nutrition". Asia Pac J Clin Nutr 2009;18:Suppl:S1-8.
Pesce M and al., Scientific Opinion on the substantiation of health claims related to live yoghurt cultures and improved lactose digestion (ID 1143, 2976) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. EFSA Journal 2010;8:1763.
Pimentel S and al., Lactose maldigestion and intolerance: pathogenesis, diagnosis and treatment. United European Gastroenterol J 2013;1:113-9.
Rizkalla SW and al., Improved lactose tolerance in healthy adults consuming yogurt containing lactobacilli and bifidobacteria. J Nutr 2004;134:2435-7.
Robles TA and al., Self-perceived lactose intolerance results in lower intakes of calcium and dairy foods and is associated with hypertension and diabetes in adults. Am J Clin Nutr 2014;99:8-15.
Scalabrin DM and al., Tolerance to low-fat yogurt containing viable Lactobacillus acidophilus cells in lactose malabsorbers. J Am Coll Nutr 1998;17:243-50.
Suarez FL and al., A comparison of symptoms after the consumption of milk or lactose-hydrolyzed milk by people with self-reported severe lactose intolerance. N Engl J Med 1995;333:1-4.
Surono IS, Bhanthin PR, Carpentier TD and al, NIH consensus development conference statement: Lactose intolerance and health. NIH Consens State Sci Statements 2010;27:1-27.
Szilagyi A and al., Adaptation, confounding and hormesis in lactose malabsorption. J Am Coll Nutr 2015;34:1-13.
Wang and al., Yogurt consumption is associated with better diet quality and metabolic profile in American women. Am J Clin Nutr 2015;101:1721-6.
Wang and al., Yogurt consumption is associated with better diet quality and metabolic profile in American women. Am J Clin Nutr 2015;101:1721-6.
WGO. World Organisation Health Day 'A global diet and lifestyle tips on how to improve digestive health. http://www.worldgastroenterology.org/UserFiles/news/WDHD/2010/2010%20Guidelines20-Spanish.pdf. Accessed 2010;1-20.
Wrigglesworth J, Cook HM, Atkinson SA, Computed models and implications for human health. Pharmacol Res 2000;42:103-10.

YOGURT NUTRITION
INITIATIVE FOR A BALANCED DIET